

无机化学学报

2019 年

第 35 卷

第 11 期

目 次

综 述

多酸基光敏剂在染料敏化太阳能电池中的应用

.....谷易桐 陈 黎 李建平 刘 丽 陈维林 刘 丁 王恩波(1905)

金属-有机框架衍生的中空碳材料及其在电化学能源存储与氧还原领域中的应用

.....刘 虎 杨东辉 王许云 韩宝航(1921)

多酸基主客体框架材料 POMs@MOFs(COFs).....李季坤 赵帅恒 胡长文(1934)

仿生矿化增强材料.....曹 含 潘海华 唐睿康(1957)

pH 敏感铈、钕和铂配合物在肿瘤靶向荧光成像和治疗中的应用(英文)

.....张思琪 高丽华 赵 华 王克志(1974)

单分子磁环最新进展及研究展望.....李晓磊(1987)

水系锌离子二次电池锌负极的研究进展.....王福慧 刘辉彪(1999)

单离子磁体及其磁弛豫动力学的研究进展.....苟晓霜 王梦梦 孟茜茜 程 鹏(2013)

论 文

一种螯合型 Fe^{2+} 探针的构筑及性质

.....李泽冉 朱健萍 于涵洋 陈韵聪 何卫江 郭子建(2031)

基于三吡啶烯烃配体的一维配位聚合物:合成、结构及光电性质

.....陈慧娴 刘春玉 周 璇 陈敏敏 倪春燕 郎建平(2038)

核壳结构 MoO_3/C 微球的制备及储锂性能

.....白玉林 王 澄 吴 越 刘于斯 马 超 蔡志鹏 王开学 陈接胜(2045)

SAPO-11 的绿色制备及在棕榈油催化加氢制备生物航煤方面的应用

.....刘思阳 白依凡 李 伟(2051)

污泥与醋糟共热解特性及碱金属迁移规律.....李强强 张阳倩 郑 颜 张毅民(2057)

双吡啶酰胺 Schiff 碱双核 Fe(II) 配合物的合成与磁性

.....张 男 苑 娟 刘美姣 金怡姝 寇会忠(2066)

纳米氧化钨的制备与热膨胀性.....孙 静 李 强 林 鲲 刘占宁 邢献然(2073)

金属单原子模型催化剂热稳定性的反应力场(ReaxFF)分子动力学研究

.....杨文琦 汪 杰 乔园园 王贵昌(2078)

生物金属有机框架 ZnBTCA 对 CO_2 的选择性吸附性能

.....才 红 吴光伟 赖佳丽 陈锦萍 陈燕飞 李 丹(2083)

4-(2-羟基-3-氯)苯基-2,2':6',2''-三联吡啶 Cu(II) 配合物的合成、结构表征及抗肿瘤活性

.....钟玉君 陈振锋 梁 宏(2089)

CYCU-3 孔内合成聚吡咯及其电导性能表征.....	周盈汐	王会锋	孙福兴	(2095)
一例夹心型混合价铋钨酸盐的合成、结构及性质				
.....	杨宗斐	牟秋水	王亚平	马鹏涛
	王敬平	牛景杨		(2101)
铈簇基纳米分子胶囊的合成、晶体结构和荧光性质(英文)				
.....	陶艳丽	陈维超	王新龙	苏忠民
				(2108)
双功能镧系化合物:合成、结构、磁性和荧光性质(英文).....	韩永芳	蔡丽珍	郭国聪	(2117)
荧光介孔二氧化硅纳米粒子的合成及药物运输(英文)				
.....	陈敏敏	耿浩然	胡金霞	张 琮
	Godfred Amfo Agyekum	张卓琦	曹希传	(2125)
全氟酰基合成后修饰 UiO-66 用于吸附有机污染物(英文)				
.....	许梦颖	宋甘霖	韩宝航	(2136)
2,6-二(1-咪唑基)萘和二羧酸构筑的金属-有机框架化合物:合成、晶体结构和荧光识别性能(英文)				
.....	刘志强	曹师虎	张 哲	武峻峰
	赵 越	孙为银		(2145)
离子受体对自由卟啉探针荧光识别性能的影响(英文)				
.....	耿 聪	郑 慧	陈玉婷	姜建壮
				(2152)
3,3'-(吡啶-3,5)二苯甲酸构筑的 Co(II)/Ni(II)配合物的合成、晶体结构及性质(英文)				
.....	韩 晓	邵志超	赵 蓓	任 宁
	孟祥茹	丁 洁	侯红卫	(2159)
两种含氮类大豆苷元衍生物对人体红细胞的抗溶血作用(英文)				
.....	刘会青	于溪璇	宋静磊	延 玺
	徐 宁	郝海军		(2168)
基于多面体低聚倍半硅氧烷体系的一种独特发光现象(英文)				
.....	张庆瑞	宋明星	邓瑞平	周 亮
	张洪杰			(2177)
水热合成具有分级结构的钨酸铋纳米花及其光催化降解四环素性能(英文)				
.....	张宇晴	曾雪玉	于 凯	刘桂芳
	曹海雷	吕 健	曹 荣	(2185)
《无机化学学报》投稿须知.....				(2192)

CHINESE JOURNAL OF INORGANIC CHEMISTRY


Vol.35

No.11

Nov. 2019

CONTENTS

Cover



Special Issue Dedicated to the 100th Anniversary of the Birth of Prof. Chen Rongti


Reviews

Application of Polyoxometalates-Based Photosensitizers in Dye-Sensitized Solar Cells

GU Yi-Tong, CHEN Li, LI Jian-Ping, LIU Li, CHEN Wei-Lin, LIU Ding, WANG En-Bo

DOI:10.11862/CJIC.2019.227

Chinese J. Inorg. Chem., **2019**,*35*(11):1905-1920




Metal-Organic Framework-Derived Hollow Carbon Materials for Electrochemical Energy Storage and Oxygen Reduction Reaction

LIU Hu, YANG Dong-Hui, WANG Xu-Yun, HAN Bao-Hang

DOI:10.11862/CJIC.2019.237

Chinese J. Inorg. Chem., **2019**,*35*(11):1921-1933




In this review, the preparation processes and electrochemical performances of hollow carbon materials derived from metal-organic frameworks are outlined, mainly including their applications in lithium-ion batteries, lithium-sulfur/selenium batteries, sodium-ion batteries, supercapacitors, and oxygen reduction reaction.

Polyoxometalate-Based Host-Guest Framework Materials POMs@MOFs(COFs)

LI Ji-Kun, ZHAO Shuai-Heng, HU Chang-Wen

DOI:10.11862/CJIC.2019.230


Chinese J. Inorg. Chem., **2019**,**35**(11):1934-1956



This review summarizes the recent research progress of the POMs@MOFs(COFs) materials. The synthesis, structures and properties of these materials have been discussed in detail. The future development trend of the POMs@MOFs(COFs) materials is also prospected.

Materials Enhanced by Biomimetic Mineralization

CAO Han, PAN Hai-Hua, TANG Rui-Kang




This review mainly introduces the natural reinforced composite materials from four types of minerals, and the artificial organic-inorganic reinforced composite materials with three kinds of matrices.

pH-Sensitive Iridium, Ruthenium and Platinum Complexes for Tumor-Specific Fluorescence Imaging and Cancer Therapy (English)

ZHANG Si-Qi, GAO Li-Hua, ZHAO Hua,
WANG Ke-Zhi

DOI:10.11862/CJIC.2019.231

Chinese J. Inorg. Chem., **2019**,**35**(11):1974-1986




Acidic tumor environment is a significant hallmark of tumor tissues and provides a powerful platform for accurate tumor diagnosis and efficient therapy. Due to excellent photophysical properties and anticancer activities, pH-sensitive metal complexes of iridium, ruthenium and platinum have the potential to become tumor imaging and therapy agents.

Single-Molecule Toroids: Recent Advances and Perspectives

LI Xiao-Lei

DOI:10.11862/CJIC.2019.246

Chinese J. Inorg. Chem., **2019**,**35**(11):1987-1998




Up-to-date overview of the emerging single-molecule toroids (SMTs) are presented here in detail to expound the influencing factors and designing strategies of SMTs and ultimately inspire the searching of SMTs with enhanced toroidal moment.

Research Progress of Zinc Anode Materials for Aqueous Zinc Ion Recharge Battery

WANG Fu-Hui, LIU Hui-Biao

DOI:10.11862/CJIC.2019.239

Chinese J. Inorg. Chem., **2019**,**35**(11):1999-2012




Zinc ion recharge battery, a new type of green environmental protection battery, the performance of zinc anode can be improved by controlling its morphology and surface modification. Herein, we reviews the development of anode materials for zinc ion batteries, including metal zinc anode, composite zinc anode and zinc alloy.

Research Progress on Single-Ion Magnets and Their Magnetic Relaxation Dynamics

GOU Xiao-Shuang, WANG Meng-Meng, MENG Xi-Xi, CHENG Peng

DOI:10.11862/CJIC.2019.232

Chinese J. Inorg. Chem., **2019**,**35**(11):2013-2030



Single-ion magnets with high performance and their magnetic relaxation mechanism are reviewed.


Articles

Construction and Fe²⁺ Sensing of a Chelation Based Fe²⁺ Probe

LI Ze-Ran, ZHU Jian-Ping, YU Han-Yang, CHEN Yun-Cong, HE Wei-Jiang, GUO Zi-Jian

DOI:10.11862/CJIC.2019.223

Chinese J. Inorg. Chem., **2019**,**35**(11):2031-2037




The probe BTPY was constructed via integrating a BODIPY fluorophore with terpyridine at α -position by a vinyl group. By combining absorption and fluorescence spectroscopy, BTPY is able to detect iron ions visually from metal ions commonly found in biological systems, and especially capable of distinguishing Fe²⁺ from Fe³⁺.

One-Dimensional Coordination Polymers Based on a Tripyridine Olefin Ligand: Synthesis, Structures and Photoelectronic Properties

CHEN Hui-Xian, LIU Chun-Yu, ZHOU Xuan, CHEN Min-Min, NI Chun-Yan, LANG Jian-Ping

DOI:10.11862/CJIC.2019.228

Chinese J. Inorg. Chem., **2019**,**35**(11):2038-2044




The reactions of a tripyridine olefin ligand (ppvppa) with Ag(I) or Ni(II) salts afforded four one-dimensional coordination polymers. Ag(I)-based complexes **1**~**3** showed good luminescent properties while no luminescence was observed for Ni(II)-based complex **4**. Complexes **1** and **4** as representative samples showed relatively prompt, rapid anodic photocurrent responses.

Preparation and Performance for Lithium Storage of Core-Shell-Structured MoO_3/C Microspheres

BAI Yu-Lin, WANG Cheng, WU Yue,
LIU Yu-Si, MA Chao, CAI Zhi-Peng,
WANG Kai-Xue, CHEN Jie-Sheng

DOI:10.11862/CJIC.2019.233

Chinese J. Inorg. Chem., **2019**,**35**(11):2045-2050




Core-shell-structured MoO_3/C microspheres were prepared through the calcination of a $\text{MoO}_3/\text{resin}$ precursor and showed excellent electrochemical lithium storage performance when used as an anode material for lithium-ion batteries.

Green Preparation of SAPO-11 and Application in the Production of Bio-aviation Fuel by Catalytic Hydrogenation of Palm Oil

LIU Si-Yang, BAI Yi-Fan, LI Wei

DOI:10.11862/CJIC.2019.242

Chinese J. Inorg. Chem., **2019**,**35**(11):2051-2056




Bio-aviation fuel can be obtained by hydroisomerization palm oil HDO product (C15~C18) through SAPO-11 with reduced template as to reducing environment pollution.

Co-pyrolysis Characteristics of Sewage Sludge and Vinegar Residue and Migration Law of Alkali Metals

LI Qiang-Qiang, ZHANG Yang-Qian,
ZHENG Yan, ZHANG Yi-Min

DOI:10.11862/CJIC.2019.252

Chinese J. Inorg. Chem., **2019**,**35**(11):2057-2065




Co-pyrolysis of sewage sludge and vinegar residue was conducted in a thermogravimetric analyzer (TGA) and vacuum fixed bed reactor to investigate the effects of interactions on the products, kinetic behavior and the alkali metals migration rules.

Synthesis and Magnetism of Binuclear $\text{Fe}(\text{II})$ Complexes Based on Bis-pyridylhydrazone Schiff Base Ligands

ZHANG Nan, YUAN Juan, LIU Mei-Jiao,
JIN Yi-Shu, KOU Hui-Zhong

DOI:10.11862/CJIC.2019.234

Chinese J. Inorg. Chem., **2019**,**35**(11):2066-2072




Compartmental bis-pyridylhydrazones coordinate to two $\text{Fe}(\text{II})$ ions, giving rise to high-spin binuclear complexes.

Synthesis and Thermal Expansion in Nanosized Hafnium Oxide

SUN Jing, LI Qiang, LIN Kun, LIU Zhan-Ning,
XING Xian-Ran

DOI:10.11862/CJIC.2019.249

Chinese J. Inorg. Chem., **2019**,**35**(11):2073-2077




Hafnium oxide nanocrystals with average size of 4 nm were prepared. The thermal expansion property were measured by using the technique of X-ray pair distribution function. Compared with bulk material, hafnium oxide nanocrystals had an enhanced anisotropy on lattice thermal expansion, which derived from the structure distortion in nanocrystals.

Thermal Stability of Single Atom Metal Catalysts: ReaxFF Molecular Dynamics Study

YANG Wen-Qi, WANG Jie, QIAO Yuan-Yuan,
WANG Gui-Chang

DOI:10.11862/CJIC.2019.251

Chinese J. Inorg. Chem., **2019**,**35**(11):2078-2082




The stability research of single atom metal catalyst (SAC) model performed by MD simulation shows that Fe₁/Fe(100) can exist stable at relatively high temperature (~500 K) while other M₁/M(111) (M=Cu, Pt, Pd et.al) models occur aggregation phenomenon.

Selective Adsorption of CO₂ by Biological Metal-Organic Framework ZnBTCA

CAI Hong, WU Guang-Wei, LAI Jia-Li,
CHEN Jin-Ping, CHEN Yan-Fei, LI Dan

DOI:10.11862/CJIC.2019.236


Chinese J. Inorg. Chem., **2019**,**35**(11):2083-2088



The unique adsorption behavior of ZnBTCA for CO₂ involving weak chemical interaction of N-active sites and molecular sieve effect.

Synthesis, Structural Characterization and Antitumor Activity of Copper(II) Complex of 4-(2-Hydroxy-3-chlorine)phenyl-2,2':6',2'-terpyridine

ZHONG Yu-Jun, CHEN Zhen-Feng,
LIANG Hong



The anti-tumor activity of a new copper(II) complex (**1**) of 4-(2-hydroxy-3-chloro)phenyl-2,2':6',2'-tribipyridine was synthesized and characterized. Its IC₅₀ value is (3.36 ± 0.43) μmol · L⁻¹ toward MGC80-3 cells. Complex **1** could significantly induce apoptosis of MGC80-3 cells.

DOI:10.11862/CJIC.2019.253


Chinese J. Inorg. Chem., **2019**,**35**(11):2089-2094

Synthesis and Conductivity of Polypyrrole in CYCU-3

ZHOU Ying-Xi, WANG Hui-Feng, SUN Fu-Xing

DOI:10.11862/CJIC.2019.238

Chinese J. Inorg. Chem., **2019**,**35**(11):2095-2100




The polymerization of pyrrole was carried out in the pores of a metal-organic framework (MOF) called CYCU-3 with one-dimensional (1D) mesoporous channels. The composite combines the properties of conductivity and porosity.

Synthesis, Structure and Properties of a Sandwich-Type Polyoxomolybdate Containing Mixed-Valence Antimony

YANG Zong-Fei, MU Qiu-Shui, WANG Ya-Ping, MA Peng-Tao, WANG Jing-Ping, NIU Jing-Yang

DOI:10.11862/CJIC.2019.245

Chinese J. Inorg. Chem., **2019**,**35**(11):2101-2107




A new type of polyoxomolybdate containing mixed-valence antimony, $[(Sb_3O_3)(SbMo_6O_{24})_2]^{11-}$, has been isolated in aqueous solution, which exhibits intriguing allochroic properties.

Synthesis, Crystal Structures and Fluorescence Properties of Two Nanosized Zr-Based Molecular Capsules (English)

TAO Yan-Li, CHEN Wei-Chao, WANG Xin-Long, SU Zhong-Min

DOI:10.11862/CJIC.2019.224

Chinese J. Inorg. Chem., **2019**,**35**(11):2108-2116




Two new nanosized Zr-based MCs constructed from trinuclear $Cp_3Zr_3(\mu_3-O)(\mu_2-OH)_3$ SBUs and linear dicarboxylate ligands have been obtained, which might be excellent candidates of blue-light-emitting diode devices.

Bi-functional Lanthanide Compounds: Synthesis, Structures, Magnetism and Fluorescence (English)

HAN Yong-Fang, CAI Li-Zhen, GUO Guo-Cong

DOI:10.11862/CJIC.2019.229

Chinese J. Inorg. Chem., **2019**,**35**(11):2117-2124




The structure, magnetism and fluorescence of two isostructural lanthanide (Ln) compounds with the *N*-succinopyridine (HL) ligand, namely $[Ln(HL)_2(H_2O)_4]Cl_3$ (Ln=Pr (**1**), Eu (**2**)), have been studied and discussed. The magnetic studies revealed the presence of antiferromagnetic interaction between Pr(III) ions for **1** but weak ferromagnetic interactions between Eu(III) ions for **2**. And **2** emitted intense red luminescence at room temperature.

Synthesis of Fluorescent Mesoporous Silica Nanoparticles and Application for Intracellular Drug Delivery (English)

CHEN Min-Min, GENG Hao-Ran, HU Jin-Xia,
ZHANG Qiong, Godfred Amfo Agyekum,
ZHANG Zhuo-Qi, CAO Xi-Chuan

DOI:10.11862/CJIC.2019.244

Chinese J. Inorg. Chem., **2019**,**35**(11):2125-2135




MSNs functionalized with fluorescein isothiocyanate (FITC), as the intracellular tracer, plays critical role in anti-cancer drug delivery application due to the potential for simultaneous intracellular tracing and treatment of diseases.

Postsynthetic Modification of UiO-66 with Perfluoroalkyl for Adsorbing Organic Pollutants (English)

XU Meng-Ying, SONG Gan-Lin, HAN Bao-Hang

DOI:10.11862/CJIC.2019.247

Chinese J. Inorg. Chem., **2019**,**35**(11):2136-2144




Two hydrophobic metal-organic frameworks were prepared by postsynthetic modification. The modified materials can effectively adsorb organic liquid and aromatic compounds dissolved in water.

Metal-Organic Frameworks with 2,6-Di(1*H*-imidazol-1-yl)naphthalene and Dicarboxylate Ligands: Synthesis, Crystal Structure and Photoluminescence Sensing Property (English)

LIU Zhi-Qiang, CAO Shi-Hu, ZHANG Zhe,
WU Jun-Feng, ZHAO Yue, SUN Wei-Yin

DOI:10.11862/CJIC.2019.225

Chinese J. Inorg. Chem., **2019**,**35**(11):2145-2151



Selective sensing acetone


Cu-MOF

Receptor Influence on Sensing Performance of Metal-Free Porphyrin Sensor (English)

GENG Cong, ZHENG Hui, CHEN Yu-Ting,
JIANG Jian-Zhuang

DOI:10.11862/CJIC.2019.235

Chinese J. Inorg. Chem., **2019**,**35**(11):2152-2158




A metal-free tetra(aryl)porphyrin chemosensor with two *N,N*-bis(2-pyridylmethyl) amino groups (DPA) was synthesized for $\text{Pb}^{2+}/\text{Cu}^{2+}$ on the basis of dual-optical signals due to combined action of the rigid tetrapyrrole macrocyclic primary receptor with the no-conjugated DPA auxiliary receptor.

Synthesis, Crystal Structure, Magnetic Properties and Catalytic Activity of Co(II)/Ni(II) Complexes with 3,3'-(Pyridine-3,5-diyl) Dibenzoic Acid (English)

HAN Xiao, SHAO Zhi-Chao, ZHAO Bei, REN Ning, DING Jie, MENG Xiang-Ru, HOU Hong-Wei

DOI:10.11862/CJIC.2019.241

Chinese J. Inorg. Chem., 2019,35(11):2159-2167




Anti-hemolysis of Human Erythrocytes by Two Daidzein Derivatives Containing Nitrogen (English)

LIU Hui-Qing, YU Mei-Xuan, SONG Jing-Lei, YAN Xi, XU Ning, HAO Hai-Jun

DOI:10.11862/CJIC.2019.240

Chinese J. Inorg. Chem., 2019,35(11):2168-2176




A Unique Luminescence Behavior Based on Polyhedral Oligomeric Silsesquioxane Compounds (English)

ZHANG Qing-Rui, SONG Ming-Xing, DENG Rui-Ping, ZHOU Liang, ZHANG Hong-Jie

DOI:10.11862/CJIC.2019.248

Chinese J. Inorg. Chem., 2019,35(11):2177-2184




The non-luminous POSS AIPPOSS (a) and OIPPOSS (b) showed bright blue luminescence after heating and refluxed in their THF solutions. It is investigated and supposed to be attributed to the formation of the POSS/solvent adducts, which modulate the electronic structures of the POSS compounds, and consequently lead to the unique luminescence behavior.

Hydrothermal Synthesis of Hierarchically Structured Flower-like Bismuth Tungstate for Photocatalytic Tetracycline Degradation (English)

ZHANG Yu-Qing, ZENG Xue-Yu, YU Kai, LIU Gui-Fang, CAO Hai-Lei, LÜ Jian, CAO Rong

DOI:10.11862/CJIC.2019.243

Chinese J. Inorg. Chem., 2019,35(11):2185-2191



The hierarchically structured flower-like bismuth tungstate is able to produce various active species toward the degradation of tetracycline antibiotics (tetracycline and oxytetracycline) under visible light irradiation.